

Product datasheet MON5159-1

**MONOSAN**<sup>®</sup>

Mouse anti E-Cadherin, clone BS38 (Monoclonal)

Clone no. BS38

MONOSAN

---

Product name	Mouse anti E-Cadherin, clone BS38 (Monoclonal)
Host	Mouse
Applications	IHC-P (1:100-1:400)
Species reactivity	human, dog, rat, mouse, pig, sheep, rabbit
Conjugate	-
Immunogen	Unknown or proprietary to MONOSAN and/or its suppliers
Isotype	IgG1
Clonality	Monoclonal
Clone number	BS38
Size	1 ml
Concentration	n/a
Format	-
Storage buffer	TRIS with 0.03% sodium azide, pH7,2
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

## Mouse anti E-Cadherin, clone BS38 (Monoclonal)

Clone no. BS38

MONOSAN

**Additional info**

E-Cadherin is a 120 kDa transmembrane glycoprotein that is localized in the adherens junctions of epithelial cells. There, it interacts with the cytoskeleton through the associated cytoplasmic catenin proteins. In addition to being a calcium-dependent adhesion molecule, E-Cadherin is also a critical regulator of epithelial junction formation. Its association with catenins is necessary for cell-cell adhesion. These E-cadherin/catenin complexes associate with cortical actin bundles at both the zonula adherens and the lateral adhesion plaques. Tyrosine phosphorylation can disrupt these complexes, leading to changes in cell adhesion properties. E-Cadherin expression is often down-regulated in highly invasive, poorly differentiated carcinomas. Increased expression of E-Cadherin in these cells reduces invasiveness. Thus, loss of expression or function of E-Cadherin appears to be an important step in tumorigenic progression. Tissue specificity: Non-neural epithelial tissues.

**References**

1. -
2. -
3. -
4. -
5. -

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES