

Mouse Anti-Hu IgG (Fc) FITC, clone EM-07 (Monoclonal)

Clone no. EM-07

MONOSAN

---

Product name	Mouse Anti-Hu IgG (Fc) FITC, clone EM-07 (Monoclonal)
Host	Mouse
Applications	FC
Species reactivity	Human
Conjugate	FITC
Immunogen	Fusion protein of human IgG Fc fragment.
Isotype	IgG2
Clonality	Monoclonal
Clone number	EM-07
Size	0.1 mg
Concentration	0.1 mg/ml
Format	Purified antibody is conjugated with fluorescein isothiocyanate (FITC) under
Storage buffer	Stabilizing phosphate buffered saline (PBS), pH 7.4, 15 mM sodium azide
Storage until expiry date	2-8°C

FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES

## Mouse Anti-Hu IgG (Fc) FITC, clone EM-07 (Monoclonal)

Clone no. EM-07

MONOSAN

**Additional info**

Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the total plasma immunoglobulins. IgG has the highest stability of blood immunoglobulins (T<sub>1/2</sub> = 21 days) and is able of placental transfer. IgG is secreted by plasma cells at a comparably high rate as other immunoglobulins.

Specificity  
Immunoglobulin G (IgG) is a 150 kDa soluble protein that serves as a major effector molecule of the humoral immune response in man. Its concentration in blood plasma of healthy individuals is approximately 10 g/l, which accounts for about 75% of the total plasma immunoglobulins. IgG has the highest stability of blood immunoglobulins (T<sub>1/2</sub> = 21 days) and is able of placental transfer. IgG is secreted by plasma cells at a comparably high rate as other immunoglobulins.

Application details  
Flow cytometry: Recommended dilution: 1-4 µg/ml.

**References**

1. -
2. -
3. -
4. -
5. -

**FOR RESEARCH USE ONLY. NOT FOR USE IN DIAGNOSTIC PROCEDURES**